REPORT DOCUMENTATION PAGE						ı
Public reporting burden for this collection of in	formation is esti	mated to average 1 hour per	response, including	AFRL-SR-A	R-TR-02-	5,
Public reporting burden for this collection of information is estimated to average 1 hour per response, including gathering and maintaining the data needed, and completing and reviewing the collection of information. Send collection of information, including suggestions for reducing this burden, to Washington Headquarters Services Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperw					0384	
1. AGENCY USE ONLY (Leave blank) 2. REPORT DATE 3. REPOR					0 00 1	7
		18 Sep 02	FIN		01 MAR 01 TO 28 FEB 02	╛
4. TITLE AND SUBTITLE (DURIP-01) PURE-BEAMS OF HIGHLY VIBRATIONALLY EXCITED MOLECULES					UNDING NUMBERS 620-01-1-0201 4/US	
6. AUTHOR(S)						١
Prof. Alec M. Wodtke					03D	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)					PERFORMING ORGANIZATION	┨
University of California, Santa Barbara					REPORT NUMBER	١
Department of Chemistry						1
Santa Barbara, CA 93106						1
						ı
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)					SPONSORING/MONITORING	\dashv
AFOSR/NL					AGENCY REPORT NUMBER	١
4015 Wilson Blvd, Room 713						١
Arlington, VA 22203-1954						1
						1
44 CUDDIEMENTADY NOTES						4
11. SUPPLEMENTARY NOTES						-
						l
						١
12a. DISTRIBUTION AVAILABILITY STATEMENT				12b.	. DISTRIBUTION CODE	٦
						1
A Co. D. 11's Dalacce Distribution Halleniand						ļ
Approve for Public Release: Distribution Unlimited						1
13. ABSTRACT (Maximum 200 wo		1			TX	
The grant was used to purchase construct and install necessary instrumentation to implement Hexapole focusing of laser						
prepared vibrationally excited molecules. It consists of three differentially pumped chamber, for: 1) state preparaton, 2)						
hexapole filtering and 3) detection. In addition two Yag-pumped dye laser systems were purchased from Spectra-Physics and						
Sirah, which allow stimulated emission pumping to prepare highly vibrationally excited states of molecules. We have also didicated another existing Yag-pumped dye laser system to the project for REMPI and LIF detection of the focused molecules						
didicated another existing 1 ag-p	umped dye	laser system to the p	roject for Kr	ENIFI and LIF d	election of the focused molecul	-3
						-
;						-
		ξ				
						١
						ł
						١
		,				
•						
44 OUR FOT TERMS					11E NUMBER OF BACES	4
14. SUBJECT TERMS					15. NUMBER OF PAGES	
					16. PRICE CODE	\dashv
17. SECURITY CLASSIFICATION OF REPORT	18. SECURI OF THIS	TY CLASSIFICATION S PAGE	19. SECURIT OF ABST		ON 20. LIMITATION OF ABSTRAC	7
Unclas		Unclas	Unclas			

Standard Form 298 (Rev. 2-89) (EG) Prescribed by ANSI Std. 239.18 Designed using Perform Pro, WHS/DIOR, Oct 94

Final Technical Report

In 2001 UCSB (for the research of Professor Alec Wodtke) was awarded a Defense University Research Instrumentation Program (DURIP) grant with the following particulars.

- 1. TITLE: "Pure-beams" of highly vibrationally excited molecules.
- 2. PRINCIPAL INVESTIGATOR: Professor Alec M. Wodtke (805) 893 8085/8552; University of California; Department of Chemistry; Santa Barbara, CA. 93106.
- 3. CURRENT DOD CONTRACT OR GRANT: Air Force Office of Scientific Research Grant Number F49620-01-1-0201.
- 4. AMOUNT: \$273,000.
- 5. DATES: March 1, 2001 Feb. 27, 2002.

The grant was used to purchase construct and install necessary instrumentation to implement

Hexapole focusing of laser prepared vibrationally excited molecules. A schematic diagram of the now completed instrument is shown in Fig. 1.

It consists of three differentially pumped chamber, for: 1) state preparation, 2) hexapole filtering and 3) detection. In addition two Yag-pumped




FIG. 1. Schematic of the beam source for "pure vibrationally excited NO": NO expands from a pulsed valve and is excited (laser beam orthogonal to figure surface) before the skimmer by SEP. The hexapole (Rod diameter--3 mm, Center-Center Rod spacing--9 mm, hexapole length--50 cm) selectively focuses the laser prepared molecules onto the target aperture. Initially, the source will be attached to a detection chamber, where REMPI and electron impact ionization detection of transmitted molecules is performed.

dye laser systems were purchased from Spectra-Physics and Sirah, which allow stimulated emission pumping to prepare highly vibrationally excited states of molecules. We have also dedicated another existing Yag-pumped dye laser system to the project for REMPI and LIF detection of the focused molecules.




Fig. shows photograph the instrument. The reflectron can be easily recognized (middle-toupper left). The source chamber is at the right. The Pump laser system can just be seen in the foreground and the

DUMP laser to the right in the background is also visible. The Probe laser is not visible behind the vacuum chamber. A large 3-D Helmholz coil allows experiments in well controlled magnetic fields.

We have now successfully focused vibrationally excited molecules through the hexapole demonstrating that our basic experimental premise. The performance of the instrument appears promising. We now routinely achieve electric fields in excess of 100 kVolts/cm. Focusing curves

have been simulated by classical trajectory calculations. We will be preparing publication of this result in the near future.

Preliminary results from this instrument have also been presented at several scientific forums.

- 1. Invited Speaker, ACS meeting Orlando Florida, April 7 2002.
- 2. Invited Speaker, Atomic and Molecular Interactions Gordon Conference, July 7 2002.
- 3. Invited Speaker "Chemistry Colloquium" Boston University, Boston Massachusetts, Sept. 17 2002
- 4. Invited Speaker "Physical Chemistry Seminar" Massachusetts Insitute of Technology, Boston Massachusetts, Sept. 17 2002
- 5. Invited Speaker "Physical Chemistry Seminar" University of Sherbrooke, Quebec Canda, Sept. 18 2002
- 6. Invited Speaker "Physical Chemistry Seminar" Emory University, Atlanta Georgia, Sept. 20 2002
- 7. Invited Speaker "Conference on Stereo-Dynamics of Chemical Reactions" Schoorl, the Netherlands Dec. $1-6\,2002$